

doi: 10.13241/j.cnki.pmb.2017.32.031

宫腹腔镜联合治疗输卵管阻塞性不孕症的疗效及在不同年龄及阻塞类型中疗效差异

陈丽融 刘常燕 李 莹 王 茜 胡又丹

(达州市中西医结合医院妇产科 四川 达州 635002)

摘要 目的: 探讨宫腹腔镜联合治疗输卵管阻塞性不孕症的临床治疗效果及在不同年龄及阻塞类型中的疗效差异。**方法:** 选取2013年2月至2015年2月间我院收治的70例输卵管性不孕症患者,按照随机数字表法分为两组,对照组($n=35$ 例)行宫腔镜引导下的输卵管疏通术,观察组($n=35$ 例)在对照组治疗方案上联合腹腔镜治疗,比较两组的临床治疗效果,同时比较宫腹腔镜联合对不同年龄段以及不同阻塞类型患者的临床治疗效果。**结果:** 观察组输卵管通畅情况优于对照组($Z=-2.207, P=0.027$);观察组输卵管再通率高于对照组[82.86% vs 60.00%],差异有统计学意义($\chi^2=4.480, P=0.034$)。两组治疗后并发症发生率无差异($P>0.05$)。观察组治疗后1年总妊娠率高于对照组 [82.86% vs 57.14%], 差异有统计学意义 ($P<0.05$); 宫内妊娠率高于对照组 [80.00% vs 51.43%], 差异有统计学意义 ($P<0.05$)。年龄 ≤ 30 岁组与年龄 >30 岁组患者输卵管再通率、总妊娠率、宫内妊娠率无差异($P>0.05$)。壶腹部或伞部阻塞患者输卵管再通率、总妊娠率、宫内妊娠率均高于间质部阻塞患者和峡部阻塞患者,差异有统计学意义 ($P<0.05$)。**结论:** 宫腹腔镜联合治疗输卵管阻塞性不孕症具有较好的疗效,且输卵管再通率和妊娠率得到明显提高,术后并发症也较少,并且对输卵管壶腹部或伞部阻塞的术后再通率效果最好。

关键词: 宫腔镜;腹腔镜;输卵管阻塞;不孕症;临床效果

中图分类号: R711.6; R711.76 **文献标识码:** A **文章编号:** 1673-6273(2017)32-6340-05

Study on Efficacy of Hysteroscopy Combined with Laparoscopy on Salpingemphraxis Infertility

CHEN Li-rong, LIU Chang-yan, LI Ying, WANG Qian, HU You-dan

(Department of Gynecology and Obstetrics, Dazhou Integrated Traditional Chinese and Western Medicine Hospital of Sichuan Province, Dazhou, Sichuan, 635002, China)

ABSTRACT Objective: To explore the efficacy of hysteroscope combined with laparoscope combined with in the treatment of salpingemphraxis infertility and differences in efficacy of different ages and types of obstruction. **Methods:** A total of 70 patients with salpingemphraxis infertility, who were admitted to Dazhou Integrated Traditional Chinese and Western Medicine Hospital of Sichuan Province from February 2013 to February 2015, were randomly divided into control group ($n=35$) and observation group ($n=35$). The control group ($n=35$) underwent dredge tubal operation guided by hysteroscope, while the observation group was received laparoscopic treatment on the treatment plan of the control group. The clinical effects of the two groups were compared, at the same time, the clinical effects of hysteroscopy combined with laparoscopy on different age groups and different types of obstruction were compared. **Results:** The patency of fallopian tube in the observation group was better than that in the control group ($Z=-2.207, P=0.027$). The fallopian tube recanalization rate in the observation group was higher than that in the control group [82.86%vs60.00%], the difference was statistically significant ($\chi^2=4.480, P=0.034$). There was no difference in the incidence of complications after treatment between the two groups ($P>0.05$). The overall pregnancy rate in the observation group 1 year after treatment was higher than that in the control group [82.86%vs57.14%], the difference was statistically significant ($P<0.05$). The intrauterine pregnancy rate in the observation group was higher than that in the control group [80.00%vs51.43%], the difference was statistically significant ($P<0.05$). There were no statistical differences in tubal recanalization rate, overall pregnancy rate, intrauterine pregnancy rate of patients between ≤ 30 years and >30 years ($P>0.05$). The fallopian tube recanalization rate, total pregnancy rate and uterine pregnancy rate in the patients with ampulla or umbrella obstruction were higher than those in interstitial obstructive patients and isthmus obstructive patients, the difference was statistically significant ($P<0.05$). **Conclusion:** Hysteroscope combined with laparoscope in the treatment of salpingemphraxis infertility has better clinical efficacy, and the recanalization rate and pregnancy rate of fallopian tube are obviously improved, with less postoperative complications and the highest recanalization rate of fallopian tube, ampulla or umbrella block.

Key words: Hysteroscope; Laparoscope; Salpingemphraxis; Infertility; Clinical efficacy

Chinese Library Classification(CLC): R711.6; R711.76 **Document code:** A

Article ID: 1673-6273(2017)32-6340-05

作者简介:陈丽融(1981-),女,本科,主治医师,从事妇科方面的研究, E-mail: mqoejf@163.com

(收稿日期:2017-05-16 接受日期:2017-06-10)

前言

不孕症育龄期妇女常见的疾病,指的是未采取避孕措施且正常性生活至少1年而未妊娠的现象,随着性传播疾病、宫内操作的增加以及性观念的开放,其发病率呈逐渐升高的趋势^[1,2]。输卵管阻塞是不孕症众多发病原因中最为常见的,其所占比例约为30%~40%^[3]。不孕症传统的治疗方法如子宫输卵管碘油造影术以及输卵管通液术虽然操作简单,费用较低,但是由于无法全面找出发病原因,且盲目性相对较大,因而难以达到满意的治疗效果^[4,5]。随着微创技术和内镜技术的不断发展和成熟,宫腹腔镜联合成为临床治疗输卵管性阻塞性不孕症的重要方式,它能够全面发现不孕的发病原因,且术后治疗效果好,患者满意度高^[6-8]。但是关于宫腹腔镜联合在不同年龄段以及不同阻塞类型不孕症患者中相关的研究相对较少,本研究对输卵管性不孕症患者采用宫腹腔镜联合治疗,并以单独腹腔镜引导下输卵管疏通术为对照组,旨在观察宫腹腔镜的临床治疗效果。

1 资料与方法

1.1 一般资料

选取2013年2月至2015年2月间我院收治的70例输卵管性不孕症患者为研究对象,纳入标准:①符合输卵管阻塞性不孕症的诊断标准^[9],并经传统宫腔碘油造影或者子宫输卵管造影检查确诊;②已婚妇女;③无手术禁忌症者;④患者均有怀孕诉求。排除标准:①男方因素所致的不孕;②肝、肾功能障碍者;③心功能不全者;④合并有其他可能影响妊娠的盆腔器质性病变;⑤合并有内分泌异常或者先天性的子宫发育不良导致的不孕。按照随机数字表法将其分为两组,观察组35例,年龄23~39岁,平均(27.9±5.6)岁;不孕持续时间2~8年,平均(3.9±1.0)年;原发性不孕7例,继发性不孕28例。对照组35例,年龄21~37岁,平均(26.5±6.7)岁;不孕持续时间2~7年,平均(3.5±1.2)年;原发性不孕4例,继发性不孕31例。两组患者的年龄构成、不孕持续时间及不孕症类型比较差异不显著($P>0.05$),具有可比性。将观察组患者按照不同年龄分为≤30岁组($n=21$ 例)和>30岁组($n=14$ 例),两组患者的不孕持续时间、不孕症类型差异无统计学意义($P>0.05$);将观察组患者按照不同阻塞部位分为间质部阻塞($n=8$ 例)、峡部阻塞($n=7$ 例)和壶腹部或伞部阻塞($n=20$ 例),不同阻塞部位患者的年龄构

成、不孕持续时间、不孕症类型比较差异不显著($P>0.05$)。

1.2 方法

观察组患者采用腹腔镜联合腹腔镜进行治疗,具体的操作分两步进行:^①采用腹腔镜对患者进行探查,即将气腹针经脐部插入,并充入二氧化碳以形成气腹,然后将腹腔镜置入其中以察看并确定患者的子宫形状、大小,输卵管以及卵巢的形状,从而制定出手术方案。^②在腹腔镜的引导下行输卵管疏通术,即将宫颈扩开后将腹腔镜置入其中,找到输卵管的开口位置后将医用导管插入,并加压注入混合药液(生理盐水20 mL、庆大霉素5 mg、地塞米松5 mg、糜蛋白酶1500 U)以将输卵管疏通,在整个手术的操作过程中动作要轻柔,防止损伤输卵管。对照组患者行单纯的腹腔镜治疗,再配合行与观察组相同的输卵管插管疏通术。两组患者术后均行常规的抗感染治疗。

1.3 评价指标

^①手术后第3个月,两组患者的输卵管复通率,以及手术后发生的并发症。其中输卵管的通畅率评价标准^[10]:①输卵管通畅:将美兰液指示液注入后输卵管顺利无回逆和阻力,或者开始有回逆和阻力,加压(40~80 kPa之间为最大推注压力)后的回逆和阻力消失;②输卵管梗阻:指示液注入后阻力大,将推注压力增加至40~80 kPa后无法注入,指示液全部回逆;③输卵管通而不畅:指示液注入后部分回逆、有一定阻力。^②术后对患者随访1年,记录两组患者的总妊娠率,包括宫内妊娠率、异位妊娠率。

1.4 统计学处理

采用SPSS16.0软件对数据录入及统计分析,一般资料的年龄等计量资料以($\bar{x}\pm s$)表示,两组独立样本的比较采用成组t检验,输卵管再通率、并发症发生率、妊娠率等计数资料以率(%)表示,比较采用 χ^2 检验或者Fisher确切概率法,设置检验标准为 $\alpha=0.05$ 。

2 结果

2.1 两组患者的输卵管复通率比较

治疗前,两组患者的输卵管通畅情况比较差异无统计学意义($Z=-0.403$, $P>0.687$),治疗后观察组患者输卵管的通畅情况优于对照组($Z=-2.207$, $P=0.027$);观察组治疗后的输卵管再通率为82.86%,对照组为60.00%,观察组输卵管再通率高于对照组,差异有统计学意义($\chi^2=4.480$, $P=0.034$)。见表1。

表1 比较两组患者治疗后的输卵管复通率[n(%)]

Table 1 Comparison of tubal recanalization rates between patients in two groups after treatment[n(%)]

Groups	n	Fallopian tube patency		Fallopian tube obstruction		Oviduct passage problem		Recanalization rate
		Before treatment	3 months after treatment	Before treatment	3 months after treatment	Before treatment	3 months after treatment	
Observation group	35	2(5.71)	31(88.57)	11(31.43)	3(8.57)	22(62.86)	1(2.86)	29(82.86)
Control group	35	4(11.43)	24(68.57)	10(28.57)	5(14.29)	21(60.00)	6(17.14)	21(60.00)

2.2 两组患者治疗后的并发症发生率比较

两组患者治疗后有不同程度的并发症发生,包括阴道出血、腹部疼痛,两组并发症发生率比较,差异无统计学意义

($P>0.05$)。见表2。

2.3 两组患者治疗后1年的妊娠率比较

观察组治疗后1年的总妊娠率为82.86%,对照组为

57.14%, 观察组总妊娠率高于对照组, 差异有统计学意义 ($P < 0.05$)。见表 3。
05); 观察组宫内妊娠率 80.00%, 高于对照组的 51.43%, 差异有

表 2 比较两组患者治疗后的并发症发生率[n(%)]

Table 2 Comparison of complications after treatment between two groups[n(%)]

Groups	n	Vaginal bleeding	Abdominal pain	Infected
Observation group	35	3(8.57)	2(5.71)	0(0.00)
Control group	35	5(14.29)	3(8.57)	0(0.00)
χ^2	-	0.565	0.215	-
P	-	0.452	0.643	-

表 3 比较两组患者治疗后 1 年的妊娠率[n(%)]

Table 3 Comparison of pregnancy rates 1 year after treatment between two groups [n(%)]

Groups	n	Total pregnancy rate	Intrauterine pregnancy rate	Ectopic pregnancy rate
Observation group	35	29(82.86)	28(80.00)	1(2.86)
Control group	35	20(57.14)	18(51.43)	2(5.71)
χ^2	-	5.510	6.341	-
P	-	0.019	0.012	1.000*

Note: * calculated by exact probability method of Fisher.

2.4 观察组不同年龄段患者的输卵管再通率、妊娠率比较 妊娠率、宫内妊娠率比较, 差异无统计学意义 ($P > 0.05$)。见表 4。
年龄 ≤ 30 岁组与年龄 > 30 岁组患者的输卵管再通率、总

表 4 比较观察组不同年龄段患者的输卵管再通率、妊娠率[n(%)]

Table 4 Comparison of fallopian tube recanalization rate and pregnancy rate in different age groups in observation group[n(%)]

Groups	n	Fallopian tube recanalization rate(n=29)	Total pregnancy rate(n=29)	Intrauterine pregnancy rate (n=28)
≤ 30 years group	21	17(80.95)	15(71.43)	18(85.71)
> 30 years group	14	12(85.71)	14(100.00)	10(71.43)
χ^2	-	0.134	-	1.071
P	-	0.714	0.061*	0.301

Note: * calculated by exact probability method of Fisher.

2.5 观察组不同阻塞部位患者的输卵管再通率、妊娠率比较 均高于间质部阻塞患者和峡部阻塞患者, 差异有统计学意义
壶腹部、伞部阻塞患者输卵管再通率、总妊娠率、宫内妊娠 ($P < 0.05$)。见表 5。

表 5 比较观察组不同阻塞部位患者的输卵管再通率、妊娠率[n(%)]

Table 5 Comparison of fallopian tube recanalization rate and pregnancy rate in different obstructive places of observation group[n(%)]

Groups	n	Fallopian tube recanalization rate(n=29)	Total pregnancy rate(n=29)	Intrauterine pregnancy rate (n=28)
Interstitial obstruction	8	4(50.00)	5(62.50)	5(62.50)
Isthmic obstruction	7	5(71.43)	4(57.14)	4(57.14)
Ampulla or umbrella obstruction	20	20(100.00)*	20(100.00)*	19(95.00)*
χ^2	-	10.862	9.731	6.629
P	-	0.004	0.008	0.036

Note: compared with the other two groups, * $P < 0.05$.

3 讨论

不孕症已成为育龄期妇女常见的生殖系统疾病,近年来,其在我国的发病率呈逐年增加的趋势,给有生育要求的女性带来严重的困扰。输卵管在女性的生殖系统中是最重要的组成部分之一,具有输送精子、摄取卵子、运送受精卵等功能,一旦输卵管发生障碍,极有可能导致女性不孕^[11-13]。已有研究证实^[14-16],输卵管阻塞是引起女性不孕症的重要原因,其病理机制为各种急、慢性输卵管炎症导致输卵管阻塞,致使精子和卵子结合受阻,最终引发输卵管阻塞性不孕症。目前,针对输卵管阻塞性不孕症的治疗方法主要有传统中医中药方案、宫腔内通液、宫腔镜下再通术的应用等^[17,18]。其中传统的中医中药起效较慢,疗程较长,是一个长期的过程;宫腔内通液治疗则常因治疗药物难以准确进入输卵管,或者无法明确输卵管阻塞是一侧还是双侧,因此治疗效果不理想,满意度较差;而单纯的宫腔镜下输卵管再通术不能较好的改善输卵管粘连,同时也有增加宫外孕的风险^[19-21]。随着微创技术的快速发展,报道显示^[22],宫腔镜与腹腔镜双“镜子”齐下治疗输卵管阻塞性不孕症,能明显提高临床治疗效果。宫腔镜可以对整个子宫进行探查,以了解患者的不孕症发生具体原因,并能据实际病情采取针对性的治疗方案,而腹腔镜则可以对腹腔内的脏器清晰的察看,手术操作者可以在腹腔镜直视下进行相关手术^[23-25]。

本研究采用宫腹腔镜联合治疗输卵管阻塞性不孕症,并以单纯宫腔镜引导下的输卵管疏通术作为参照。结果显示,治疗后观察组患者输卵管的通常情况优于对照组,并且观察组输卵管再通率为82.86%,高于对照组的60.00%。结果提示宫腹腔镜联合能明显提高输卵管阻塞性不孕症患者的临床疗效,促进输卵管的再通,从而为下一步的妊娠提供条件。可能与宫腹腔镜联合可以优势互补有关^[26-28];宫腔镜可以对输卵管的子宫腔开口进行探查,并将通液管准确插入输卵管口,而在插管过程中,也会对输卵管气道疏通的作用,同时,在加压通液的过程中,也可使炎症渗出物、输卵管痉挛等导致的栓子堵塞得到有效治疗;腹腔镜直视下可以对输卵管的通畅程度、以及梗阻的部位及其周围粘连情况等探查,并针对具体情况采取相应的手术方案进行治疗。此外,手术后,两组患者均发生不同程度的并发症,包括阴道出血、腹部疼痛等,但是患者均能耐受,对症处理后并发症得到缓解,两组患者并发症发生率无明显差异,说明输卵管阻塞性不孕症采用宫腹腔镜联合治疗的安全性高,临床有重要的参考价值。术后随访1年,观察组总妊娠率为82.86%,高于对照组的57.14%,观察组宫内妊娠率为80.00%,高于对照组的51.43%。结果进一步说明宫腹腔镜联合可以提高患者的妊娠率,与有关研究结果一致^[29]。

对宫腹腔镜治疗的患者按照年龄分组发现,年龄 ≤ 30 岁组与年龄 >30 岁组患者的输卵管再通率、总妊娠率、宫内妊娠率无明显差异,说明宫腹腔镜联合治疗输卵管阻塞性不孕症的疗效与年龄无关,可能与本研究纳入的样本量相对较小有关,应在以后的研究中扩大样本量加以证实。结果还发现,输卵管壶腹部、伞部阻塞的患者其输卵管再通率、总妊娠率、宫内妊娠率均高于间质部阻塞和峡部阻塞患者,可能是因为宫腹腔镜联合下,针对输卵管壶腹部、伞部阻塞的患者,在腹腔镜直视下,

通过盆腔粘连松解可以使子宫、卵巢以及输卵管的解剖位置得到有效恢复,而在宫腔镜下可进一步行伞端粘连和积水扩张术和造口术,而对于间质部阻塞以及峡部阻塞的患者,宫腹腔镜治疗的效果相对较差^[30]。因此,建议临床上应重点关注间质部阻塞以及峡部阻塞患者,对于治疗效果不佳者,应早期着手准备实施体外受精-胚胎移植手术,以帮助患者早日怀孕。

综上所述,宫腹腔镜联合可以明显发挥各自优点,起到优势互补的作用,两者联合可以明显提高输卵管阻塞性不孕症患者的临床疗效,提高再通率,帮助患者早日妊娠,且手术安全,对输卵管壶腹部、伞部阻塞的术后再通率效果最好,临床有重要参考价值。

参考文献(References)

- [1] Nam JK. Social Perception of Infertility and Its Treatment in Late Medieval Italy: Margherita Datini, an Italian Merchant's Wife [J]. *Uisahak*, 2016, 25(3): 519-556
- [2] 吴芳,刘洋,李小刚,等.不孕症患者焦虑及影响因素[J]. *中南大学学报(医学版)*, 2015, 40(9): 1023-1028
Wu Fang, Liu Yang, Li Xiao-gang, et al. Anxiety status and influential factors in patients with infertility [J]. *Journal of Central South University(Medical Science)*, 2015, 40(9): 1023-1028
- [3] 陈慧佳,甘晓玲,胡丽娜,等.不孕症患者心理状态及心率变异性的调查分析[J]. *实用妇产科杂志*, 2016, 32(5): 346-349
Chen Hui-jia, Gan Xiao-ling, Hu Li-na, et al. Investigation and Analysis of Psychological Status and Heart Rate Variability in Patient with Infertility [J]. *Journal of Practical Obstetrics and Gynecology*, 2016, 32(5): 346-349
- [4] Sun N, Wei L, Chen D, et al. Clinical observation of fallopian tube obstruction recanalization by ozone [J]. *Pak J Med Sci*, 2017, 33(2): 290-294
- [5] Sharma JB, Sneha J, Singh UB, et al. Comparative Study of Laparoscopic Abdominopelvic and Fallopian Tube Findings Before and After Antitubercular Therapy in Female Genital Tuberculosis With Infertility[J]. *J Minim Invasive Gynecol*, 2016, 23(2): 215-222
- [6] Parry JP, Riche D, Aldred J, et al. Proximal Tubal Patency Demonstrated Through Air Infusion During Flexible Office Hysteroscopy Is Predictive of Whole Tubal Patency [J]. *J Minim Invasive Gynecol*, 2017, 24(4): 646-652
- [7] De Cicco S, Tagliaferri V, Selvaggi L, et al. Expectant management may reduce overtreatment in women affected by unexplained infertility confirmed by diagnostic laparoscopy [J]. *Arch Gynecol Obstet*, 2017, 295(2): 427-433
- [8] Kabadi YM, Harsha B. Hysterolaparoscopy in the Evaluation and Management of Female Infertility [J]. *J Obstet Gynaecol India*, 2016, 66(Suppl 1): 478-481
- [9] 阳子兑,梯.中西医结合治疗输卵管阻塞性不孕症临床观察[J]. *中医药导报*, 2011, 17(7): 34-35
Yang Zhi-dui-di. Clinical observation on treatment of infertility caused by oviduct obstruction by combination of TCM and Western Medicine [J]. *Guiding Journal of Traditional Chinese Medicine and Pharmacology*, 2011, 17 (7): 34-35
- [10] 陈秋兰,袁品丹,李坚,等.应用阴道二维超声和三维子宫输卵管超声对评价输卵管通畅性的比较 [J]. *中国医药指南*, 2013, (15):

- 662-663
- Chen Qiu-lan, Qiu Pin-dan, Li Jian, et al. Application of transvaginal ultrasound and three-dimensional ultrasound of the uterus and fallopian tube patency of fallopian tube evaluation comparing [J]. Guide of China Medicine, 2013, 11(15): 662-663
- [11] Chen ZJ, Shi Y, Sun Y, et al. Fresh versus Frozen Embryos for Infertility in the Polycystic Ovary Syndrome [J]. N Engl J Med, 2016, 375(6): 523-533
- [12] Taeibi M, Gandomani SJ, Nilforoushan P, et al. Association between infertility factors and non-physical partner abuse in infertile couples [J]. Iran J Nurs Midwifery Res, 2016, 21(4): 368-371
- [13] Briceag I, Costache A, Purcarea VL, et al. Fallopian tubes--literature review of anatomy and etiology in female infertility [J]. J Med Life, 2015, 8(2): 129-131
- [14] 李凤玲, 谭慧珍, 何浏铭, 等. 宫腔镜联合腹腔镜在女性不孕症诊治中的应用[J]. 现代生物医学进展, 2014, 14(5): 937-939
- Li Feng-ling, Tan Hui-zhen, He Liu-ming, et al. Application of hysteroscopy combined with laparoscopy in the diagnosis and treatment of female infertility [J]. Progress in Modern Biomedicine, 2014, 14(5): 937-939
- [15] Nigam A, Saxena P, Mishra A. Comparison of Hysterosalpingography and Combined Laparohysteroscopy for the Evaluation of Primary Infertility[J]. Kathmandu Univ Med J (KUMJ), 2015, 13(52): 281-285
- [16] Sharma JB, Sneha J, Singh UB, et al. Effect of antitubercular treatment on ovarian function in female genital tuberculosis with infertility[J]. J Hum Reprod Sci, 2016, 9(3): 145-150
- [17] Shokeir T, Ebrahim M, El-Mogy H. Hysteroscopic-guided local endometrial injury does not improve natural cycle pregnancy rate in women with unexplained infertility: Randomized controlled trial[J]. J Obstet Gynaecol Res, 2016, 2(11): 1553-1557
- [18] 李丽美, 孙维峰, 程琦, 等. 输卵管阻塞性不孕症的中医药治疗研究新进展[J]. 现代中西医结合杂志, 2012, 21(17): 1921-1924
- Li Li-mei, Sun Wei-feng, Cheng Qi, et al. New progress in the study of TCM treatment of Tubal Obstructive Infertility[J]. Modern Journal of Integrated Traditional Chinese and Western Medicine, 2012, 21(17): 1921-1924
- [19] Di Spiezio Sardo A, Di Carlo C, Minozzi S, et al. Efficacy of hysteroscopy in improving reproductive outcomes of infertile couples: a systematic review and meta-analysis [J]. Hum Reprod Update, 2016, 22(4): 479-496
- [20] Bosteels J, Weyers S, Kasius J, et al. Anti-adhesion therapy following operative hysteroscopy for treatment of female subfertility [J]. Cochrane Database Syst Rev, 2015, (11): CD011110
- [21] Campo R, Meier R, Dhont N, et al. Implementation of hysteroscopy in an infertility clinic: The one-stop uterine diagnosis and treatment [J]. Facts Views Vis Obgyn, 2014, 6(4): 235-239
- [22] Puri S, Jain D, Puri S, et al. Laparohysteroscopy in female infertility: A diagnostic cum therapeutic tool in Indian setting [J]. Int J Appl Basic Med Res, 2015, 5(1): 46-48
- [23] Siam S, Soliman BS. Combined laparoscopy and hysteroscopy for the detection of female genital system anomalies results of 3, 811 infertile women[J]. J Reprod Med, 2014, 59(11-12): 542-546
- [24] Tofoski G, Antovska V. Influence of hysteroscopic metroplasty on reproductive outcome in patients with infertility and recurrent pregnancy loss[J]. Pril (Makedon Akad Nauk Umet Odd Med Nauki), 2014, 35(2): 95-103
- [25] Hu C, Chen Z, Hou H, et al. Infertility evaluation via laparoscopy and hysteroscopy after conservative treatment for tubal pregnancy[J]. Int J Clin Exp Med, 2014, 7(10): 3556-3561
- [26] 邓琳, 何玉琴, 罗志莲, 等. 宫腔腹腔镜联合诊治输卵管阻塞性不孕的临床分析[J]. 中国妇幼保健研究, 2014, 25(2): 254-257
- Deng Lin, He Yu-qin, Luo Zhi-lian, et al. Clinical analysis of hysteroscopy combined with laparoscopy in diagnosis and treatment of fallopian tube obstructive infertility[J]. Chinese Journal of Woman and Child Health Research, 2014, 25(2): 254-257
- [27] Firmal P, Yadav R, Agrawal S. A prospective study to evaluate the role of laparohysteroscopy in unexplained infertility [J]. J Obstet Gynaecol, 2015, 35(4): 386-388
- [28] Zhang E, Zhang Y, Fang L, et al. Combined hysterolaparoscopy for the diagnosis of female infertility: a retrospective study of 132 patients in china[J]. Mater Sociomed, 2014, 26(3): 156-157
- [29] Ahmadi F, Rashidy Z, Haghighi H, et al. Uterine cavity assessment in infertile women: Sensitivity and specificity of three-dimensional Hysterosonography versus Hysteroscopy[J]. Iran J Reprod Med, 2013, 11(12): 977-982
- [30] 杨宝华, 周贤梅, 徐军, 等. 输卵管阻塞性不孕症宫、腹腔镜术后治疗方法探讨[J]. 生殖与避孕, 2014, 34(6): 519-522
- Yang Bao-hua, Zhou Xian-mei, Xu Jun, et al. Discussion about the Follow-up Treatment on Salpinx Obstructive Infertility Postoperation of Hysteroscopy Combined Laparoscopy [J]. Reproduction and Contraception, 2014, 34(6): 519-522
-
- (上接第 6327 页)
- [26] McClure C, Mcpeak M B, Youssef D, et al. Stat3 and C/EBP β synergize to induce miR-21 and miR-181b expression during sepsis [J]. Immunology & Cell Biology, 2016, 95(1): 42
- [27] 武宇辉, 李成荣, 何颜霞, 等. 脓毒症患儿血浆微 RNA 表达及其与炎症细胞因子的相关性[J]. 中华儿科杂志, 2014, 52(1): 28-33
- Wu Yu-hui, Li Cheng-rong, He Yan-xia, et al. Relationship between expression of microRNA and inflammatory cytokines plasma level in pediatric patients with sepsis [J]. Chinese Journal of Pediatrics, 2014, 52(1): 28-33
- [28] Wang J F, Yu M L, Yu G, et al. Serum miR-146a and miR-223 as potential new biomarkers for sepsis [J]. Biochemical & Biophysical Research Communications, 2010, 394(1): 184-188
- [29] Hu X, Xu Z, Liu P, et al. IL-18 and miR-130a in severe sepsis patients with thrombocytopenia [J]. American Journal of Emergency Medicine, 2016, 10(Issue 1): 313