doi: 10.13241/j.cnki.pmb.2018.10.033

1.5T 磁共振不同序列用于检出妇科肿瘤盆腔淋巴结转移的比较*

周德伟 | 向莉娟 | 刘 羽 | 程博超 2 张 静 |

(1四川省资阳市第一人民医院新区放射科 四川 资阳 641300;2 华西妇儿医院放射科 四川 成都 610000)

摘要 目的:比较 1.5T 磁共振(MRI)不同序列条件下对妇科肿瘤盆腔淋巴结转移的检出情况,探讨最佳检出序列条件。方法:选取 2015 年 5 月 -2017 年 5 月初诊为卵巢癌、宫颈癌、子宫内膜癌的患者 78 例作为研究对象,均行盆腔淋巴结清扫术。所有患者术前 均行 T1 加权序列(T1WI)、T2 加权序列(T2WI)、增强扫描(T1WI+C)、弥散加权成像(DWI)检查,记录每个序列检查条件下检出的盆 腔转移淋巴结个数及分布。以病理结果作为判断的 " 金标准 " 进行对比。结果:对比病理检查结果,应用 DWI 序列对妇科肿瘤盆 腔淋巴结转移的检出率(95.8%)显著高于 T2WI-MRI 序列(85.8%)和 T1WI-MRI 序列(75.0%)(P<0.05);DWI 序列与 T1WI+C 序列 (90.8%)相比差异无统计学意义(P>0.05);DWI 阈值法与 DWI 短径法相比,淋巴结的检出率差异无统计学意义(P>0.05)。结论:应用 1.5T 磁共振检查妇科肿瘤盆腔淋巴节转移时,采用 DWI 序列扫描对于转移淋巴结的具有较高的检出率,显著优于其他序列扫描;在进行阳性淋巴结判断中,ADC 阈值法和短径法均可选用。

关键词:1.5T 磁共振;妇科肿瘤;淋巴结转移

中图分类号: R737.3 文献标识码: A 文章编号: 1673-6273(2018) 10-1965-04

Comparison of Different Scan Programs of 1.5T MRI for the Detection of Pelvic Lymph Node Metastasis in Gynecological Tumor*

ZHOU De-wei', XIANG Li-juan', LIU Yu^{1/2}, CHENG Bo-chao², ZHANG Jing¹

(1 Radiology Department, Ziyang People's Hospital, Ziyang, Sichuan, 641300, China;

2 Radiology Department, Huaxi maternity hospital, Chengdu, Sichuan, 610000, China)

ABSTRACT Objective: To compare different sequences of 1.5T MRI for the detection of gynecological tumor pelvic lymph node metastasis, and discuss the optimal detection sequence. **Methods:** 78 cases of patients who were diagnosed with ovarian cancer, cervical cancer, endometrial cancer with pelvic lymph node metastasis in gynaecology of our hospital were selected from May 2015 to May 2017, they were performed the pelvic lymph node dissection. Before the operation, every patient was examined by conventional T1 weighted imaging (T1WI), T2 weighted imaging (T2WI), T1WI contrast (T1WI + C), diffusion weighted imaging (DWI) scan program, and their lymphatic metastasis team location and number were recorded and summed. The positive ratios of lymphatic node at different scan program with the pathological results, and analyzed the two positive DWI criteria were compared. **Results:** Compared with the pathological results, and analyzed the two positive DWI criteria were compared. **Results:** Compared with the pathological results, and analyzed the two positive atio software the higher ratio at DWI program (50 teams 115 nodes) than T1WI-MRI (75.0%) and T2WI-MRI (85.8%), the positive ratio was 95.8%. But there was no significant difference between DWI sequence and T1WI+C sequence (90.8%) (P>0.05), and the detection rate of lymph nodes showed no significant difference between the DWI threshold method and DWI short diameter method (P>0.05). **Conclusion:** DWI scan program was superior to conventional T1WI and T2WI scan at 1.5T MRT. As for positive ratio criteria, both short calibration criteria and apparent diffusion coefficient (ADC) value criteria were reasonable.

Key words: 1.5T MRI; Gynecological tumor; Lymph node metastasis Chinese Library Classification(CLC): R737.3 Document code: A Article ID: 1673-6273(2018)10-1965-04

前言

某些妇科肿瘤如宫颈癌、卵巢癌等极易发生盆腔淋巴结转移,盆腔淋巴结转移是影响妇科肿瘤预后的重要因素之一^[14]。因此,在进行妇科肿瘤临床诊疗过程中,早期发现并且准确诊断转移淋巴结对制定合理的淋巴结手术清扫计划及改善患者

预后情况至关重要。当前,主要依靠常规的 CT 或 MRI 进行相关淋巴结异常的检测,并以淋巴结短径大于 1 cm 视为转移淋巴结,但该判断标准敏感度较低,尤其是对于直径小于 1 cm 的转移淋巴结假阴性率较高^[56]。目前为止,磁共振弥散加权成像(diffusion weighted imaging, DWI)是唯一能够在活体组织中进行水分子弥散测量的方法,能够感应水分子的微观扩散运动,

^{*}基金项目:四川省医学科研青年创新课题(Q15039)

作者简介:周德伟(1984-),男,本科,主治医师,研究方向:妇儿影像,E-mail: zhoudewei_1984@papmedhos.club

[△] 通讯作者:刘羽(1982-),本科,主治医师,研究方向:影像诊断,E-mail: liuyu_1982@papmedhos.club

⁽收稿日期:2017-11-14 接受日期:2017-12-08)

可以从细胞或分子水平研究各种疾病的病理状态,并能进行定性诊断^[7]。本研究拟探讨 1.5T 磁共振不同序列对于妇科肿瘤淋巴结转移的检出率,以期为盆腔淋巴结清扫术前诊断提供理论依据,现将研究结果报道如下。

1 材料与方法

1.1 研究对象

收集 2015 年 5 月 -2017 年 5 月于四川省资阳市第一人民 医院放射科进行相关妇科检查疑为恶性肿瘤伴有盆腔淋巴结 转移的患者 78 例,包括宫颈癌 34 例、卵巢癌 24 例、子宫内膜 癌 20 例。其中,绝经期患者 41 例,年龄 31~65 岁(50± 4.7)岁。 病例入选标准:0 患者在四川省资阳市第一人民医院进行手术 治疗并行盆腔淋巴结清扫术;0 术前 2 周内在放射科进行 MRI 检查(1.5T 磁共振);0 患者在磁共振检查与接收手术前未进行 任何放化疗治疗;0 患者病例资料及病理结果等资料齐全。经 过筛选后最终 78 例患者入组。

1.2 检查方法

磁共振检查仪器采用德国西门子公司 1.5T 磁共振仪 (MagnetomAvanto 1.5T)。78 例患者在术前均实施 T1 加权序列 (T1WI)、T2 加权序列(T2WI)、增强扫描(T1WI+C)、弥散加权成 像(DWI)检查。其中轴位 T1W1 扫描序列的参数设置为:TR600 ms、TE8.9 ms、层距 0.5 mm、层厚 2 mm、FOV30 cm、NEX4、矩 阵 320×192;周围 T2WI 脂肪抑制扫描序列参数设置为: TR3000 ms、TE119.9 ms、带宽 41.7 kHz、层距 0.5 mm、层厚 2 mm、视野 30 cn、激励次数 4 次、矩阵 288×256;增强扫描采用 扎喷酸葡胺(国药准字 H19991368,上海旭东海普药业有限公 司),给药剂量 0.2 mL/kg 体重,扫描参数设置同 T1WI。DWI 扫 描参数设置为 b=600 s/mm², TR=4000 ms, TE=57.4 ms, FOV=30 cm, NEX=4,带宽 =250 kHz, 层距 =0.5 mm,矩阵 128×128。采 用仪器处理软件做出表观弥散系数(ADC)图,并与 T2WI 扫描 图融合,选取所需淋巴结,得到 ADC 值及短径值。

1.3 治疗方法

针对患者的肿瘤类型选择相适应的根治性手术及盆腔淋 巴结清扫术。术中记录清扫淋巴结的部位及个数,并对每个淋 巴结进行病理确诊。

1.4 观察指标

记录并对比每个序列检查条件下盆腔淋巴结个数,术中记 录实际盆腔淋巴结清扫术个数,包括双侧髂内、髂外、髂总、闭 孔、闭孔窝、腹股沟淋巴共12组淋巴结,同时计算不同序列条 件下妇科肿瘤盆腔淋巴结转移的检出率。结合 DWI 序列扫描 图,分别以 ADC 阈值及淋巴结短径为判断标准,分析阳性检出 率。在四中扫描序列的扫描图像上,查找双侧髂总、髂外、闭孔、 髂内、闭孔窝、腹股沟深淋巴结,将短径大于1.0 cm 的标记为 转移淋巴结。

1.5 统计学分析

分别对每个患者进行四种序列的扫描(T1WI、T2WI、 T1WI+C、DWI),统计检出的淋巴结个数、短径、ADC值等,计算 4种扫描序列的阳性检出率、短径平均值等。计量资料用均数 加减标准差表示(x±s),多组间比较采用方差分析。计数资料采 用例数(n)表示,组间比较采用卡方检验,以 P<0.05 为差异有统 计学意义。

2 结果

2.1 MRI 扫描及病检检出结果

T1WI-MRI 扫描结果:对 78 例患者的 12× 78 组淋巴结进 行扫描后,共 21 例患者 35 组,合计 90 个淋巴结以短径判断标 准诊断为转移性淋巴结,其平均短径为(11.2±3.5)mm。 T2WI-MRI 扫描结果:21 例患者共 45 组,合计 103 个淋巴结以 短径判断标准诊断为转移性淋巴结,其平均短径为(9.3±2.2) mm。T1WI 增强扫描结果:21 例患者共 45 组,合计 109 个淋巴 结以短径判断标准诊断为转移性淋巴结,其平均短径为(11.1± 2.3)mm。结果见表 1。

经 DWI 扫描检测所有患者的淋巴结后,共有 21 例患者合 计 50 组淋巴结,115 个淋巴结经过阈值标准判断为肿瘤转移 淋巴结;扫描结果经短径判断标准共 21 例患者,合计 50 组淋 巴结,108 个淋巴结判断为肿瘤转移淋巴结,短径值(8.8± 2.9) mm。结果见表 1。

病检结果:对所有淋巴结进行病理检查中,21 例患者合计 50 组,120 个经病检确诊为转移淋巴结。结果见表 1。

Table 1 The positive lymph nodes were detected by biopsy and different sequential scalining									
Inspection method	n	Cases (groups/cas- es)	The common iliac group (group/cases)	The external iliac group (group/cases)	Obturator group (group/case)	Internal iliac group (group/case)	Obturator fossa group (group/case)	Deep inguinal group (group/case)	Average short diameter (mm)
T1WI-MRI	21	35/90	7/7	12/43	7/15	7/15	0/0	2/10	11.2± 3.5
T2WI-MRI	21	45/103	1/14	18/48	8/18	8/15	5/3	5/5	9.3± 2.2
T1WI+C	21	45/109	7/14	18/45	7/18	7/22	3/0	3/10	11.1± 2.3
Pathological examination	21	50/120	8/16	19/54	9/20	8/19	3/6	3/5	-
DWI(threshold value)	21	50/115	8/15	19/50	9/19	8/17	3/6	3/8	-
DWI(minor axis)	21	50/108	8/13	19/48	9/18	8/16	3/6	3/7	8.8± 2.9

表 1 病检及不同序列扫描阳性淋巴结检出结果

2.2 不同扫描序列下转移性淋巴结检出结果与病理结果的比较

不同扫描序列下,以 DWI 扫描(阈值法)检出阳性淋巴结最 多(50 组 115 个),与病理检查结果相比(50 组 120 个),其阳性 检出率为 95.8%(115/120);与病理检查结果比较,T1WI-MRI 序 列扫描检出阳性淋巴结 35 组共 90 个,阳性检出率为 75.0% (90/120);T2WI-MRI 序列扫描检出阳性淋巴结 45 组共 103 个,阳性检出率为 85.8%;T1WI+C 序列增强扫描检出阳性淋巴 结 45 组共 109 个,阳性检出率为 90.8%(109/120)。

T1WI-MRI 扫描与 DWI-MRI 扫描相比, 阳性检出率差异 有统计学意义 (P<0.001);T2WI-MRI 扫描与 DWI-MRI 扫描结 果相比,阳性检出率差异有统计学意义(P=0.007);T1WI+C 增 强扫描检出结果与 DWI-MRI 扫描结果相比, 阳性检出率差异 无计学意义(P=0.121)

2.3 DWI-MRI 序列检查不同判断标准阳性检出结果比较

采用 DWI-MRI 序列检查转移淋巴结,与病理检查结果相比,阈值判断标准检出 50 组共 115 个转移淋巴结,检出率为 95.8%;短径判断标准,检出 50 组共 108 个转移淋巴结,检出率 为 90%。两种判断标准相比差异无统计学意义(P=0.078)。

3 讨论

多种妇科盆腔恶性肿瘤均可累及女性生殖系统,主要包括 卵巢癌、宫颈内膜癌、宫颈癌等^[89]。不同的肿瘤常常累及女性生 殖系统的不同部位,相对应的淋巴结的转移也会出现显著差 别。卵巢癌主要累及附件,癌细胞主要由2种途径转移,分别是 顺卵巢血管上行至腹主动脉及腰部淋巴结和炎阔韧带下行至 髂内外及髂总淋巴结,另外存在极少数经圆韧带转移至腹股沟 淋巴结^[10-12]。宫颈癌主要累及宫颈,以髂外淋巴结转移多见^[13-15]。 子宫内膜癌主要累及宫颈及宫体,以髂内、闭孔、髂外为主,并 逐步累及髂总淋巴结,其中以髂外组淋巴结最易受累^[16]。因此, 本研究主要选取包括卵巢癌、宫颈内膜癌、宫颈癌等3种女性 盆腔恶性肿瘤进行研究。

由于本研究旨在比较 1.5 T 磁共振不同序列下对于妇科肿 瘤盆腔淋巴结转移检出率的差别,同时因病例较少,所以研究 中不涉及具体肿瘤淋巴结转移的规律及特点。本研究结果表明 DWI 扫描的转移淋巴结检出率最高,与病检结果相比阈值法 达到 95.8%,与 T1WI(75.0%)、T2WI(85.8%)具有显著的差异。 DWI 扫描图片分析中,以淋巴结短径为判断标准时,淋巴结阳 性检出率与阈值法和 T1WI+C 扫描未表现出明显差异,在检测 工作过程中,对于这三种方法均可使用。其原因在于 T1WI、 T2WI和 T1WI+C 主要根据扫描图片的形态学特征进行判断, 这种诊断方法对于结果的判断一方面受扫描图像的空间分辨 率等因素影响较大。另一方面,转移淋巴结与周围组织或器官 的信号差异也会对扫描的检出率造成影响,例如 T2WI 相对于 T1WI 更容易检出病变灶含水量的差异,而 T1WI+C 相比前两 者来说对于转移病灶的信号变化更加突出,所以检出率相比前 两者高。本研究结果也证明了这一点:T1WI+C的对转移淋巴 结的检出率较 T1WI 和 T2WI 的检出率高。肿瘤细胞核浆比 大,细胞内外间隙较小,排列紧密^[17,18],因而细胞中的水分子含 量较高,分子扩散受到一定的限制,当发生淋巴结转移时,其与 正常的淋巴细胞相比,将呈现不同的信号强度^[19-21]。而在 DWI 扫描成像中,由于突出了水成分高信号的特性,因而在理论上, DWI 扫描具有较高阳性检出率^[223]。在本研究中,转移淋巴结由 于水成分较高,经 DWI 扫描后将呈现出较高的信号,ADC 峰 值也出现降低,同时在病灶周围较低的背景信号衬托下,更加 容易观察,在分析过程中可以清晰的看到阳性淋巴结的边界, 这也使得短径判断阳性成为可能。同时,可以对呈现出的高信 号区域信号轻度进行测定,通过信号强度判断阳性淋巴结。

既往提出的关于妇科肿瘤盆腔淋巴结转移的判断标准,如淋巴结直径为5mm、6mm、8mm和1cm的诊断标准^[2420],灵 敏度和特异度均未得到临床工作者及学术界的广泛认可。因此,学术界有学者推荐使用最小ADC值小于0.983×10⁻³mm²/s 为判断标准^[27]。本研究参考了这一判断标准并与1cm判断标 准进行了比较,结果显示两种判断标准在淋巴结转移检出率方 面的差异缺乏统计学意义。有研究显示超过50%的子宫内膜癌 盆腔转移淋巴结的直径在1cm以下^[28-30],分析原因可能是由于 在扫描检查过程中,对于直径1cm以内的转移淋巴结体积过 小,ADC图上未出现信号增强或者难以测量,这也提示在今后 临床工作或者学术研究中,应当对直径小于1cm的转移淋巴 结进行重点关注,提出更加切合临床实际的转移淋巴结判断标 准,需要进一步进行相关研究。

综上所述,在妇科肿瘤盆腔淋巴结转移的检测中应用 1.5T 磁共振能有效提升检出率,不同序列扫描下对转移淋巴结的检 出率以 DWI 较高,与病检结果相比差异不显著,具有较好的应 用效果,并且在临床应用中避免了造影剂的使用,具有一定的 临床应用价值。但在临床工作中,由于 DWI 相对于增强扫描而 言并未显著提高妇科肿瘤盆腔转移淋巴结的检出率,因此对于 已经进行增强扫描的患者,结合成本效益分析可以不加 DWI 序列扫描。

参考文献(References)

- [1] Liang P, Hong J H, Liu G, et al. Myofibroblasts correlate with lymphatic microvessel density and lymph node metastasis in early-stage invasive colorectal carcinoma [J]. Anticancer Research, 2016, 25(4): 2705-2712
- [2] De B R, Takes R P, Castelijns J A, et al. Advances in diagnostic modalities to detect occult lymph node metastases in head and neck squamous cell carcinoma[J]. Head Neck, 2016, 37(12): 1829-1839
- [3] Nottegar A, Veronese N, Senthil M, et al. Extra-nodal extension of sentinel lymph node metastasis is a marker of poor prognosis in breast cancer patients: A systematic review and an exploratory meta-analysis
 [J]. European Journal of Surgical Oncology the Journal of the European Society of Surgical Oncology & the British Association of Surgical Oncology, 2016, 42(7): 919
- [4] Zhao J, Ren G, Cai R, et al. Accuracy of multidetector-row CT in diagnosing lymph node metastasis in patients with early gastric cancer[J]. Journal of Diagnostics Concepts & Practice, 2016, 31(5): 125-129
- [5] Yue Z, Song G, Yong H, et al. A Pilot Study of 18 F-Alfatide PET/CT Imaging for Detecting Lymph Node Metastases in Patients with Non-Small Cell Lung Cancer[J]. Scientific Reports, 2017, 7(1): 2877
- [6] Jin X, Liang N, Wang M, et al. Integrin Imaging with (99m) Tc-3PRGD2 SPECT/CT Shows High Specificity in the Diagnosis of Lymph Node Metastasis from Non-Small Cell Lung Cancer [J]. Radi-

ology, 2016, 281(3): 150813

- [7] Wang H, Zhou Z, Li Y, et al. Comparison of machine learning methods for classifying mediastinal lymph node metastasis of non-small cell lung cancer from18F-FDG PET/CT images[J]. Ejnmmi Research, 2017, 7(1): 11
- [8] Chen J, Cui Y U, Liu L, et al. CCR7 as a predictive biomarker associated with computed tomography for the diagnosis of lymph node metastasis in bladder carcinoma [J]. Oncology Letters, 2016, 11(1): 735
- [9] Suh C H, Baek J H, Choi Y J, et al. Performance of CT in the Preoperative Diagnosis of Cervical Lymph Node Metastasis in Patients with Papillary Thyroid Cancer: A Systematic Review and Meta-Analysis [J]. Ajnr American Journal of Neuroradiology, 2016, 38(1)
- [10] Wu Ruo-yu, Jia Xiang, Cao Yi-meng. Research Progress on Exosome in the Diagnosis and Treatment of Malignant Gynecologic Tumors[J]. Journal Of International Obstetrics And Gynecology, 2017, 44 (3): 276-279
- [11] Wang G M, Liu D F, Xu Y P, et al. PET/CT imaging in diagnosing lymph node metastasis of esophageal carcinoma and its comparison with pathological findings [J]. European Review for Medical & Pharmacological Sciences, 2016, 20(8): 1495
- [12] Mokhtar M, Tadokoro Y, Nakagawa M, et al. Triple assessment of sentinel lymph node metastasis in early breast cancer using preoperative CTLG, intraoperative fluorescence navigation and OSNA [J]. Breast Cancer, 2016, 23(2): 1-9
- [13] Derlin T, Schumacher U, Bengel F M. Nonneoplastic Neuroma After Radical Prostatectomy Is Not a Mimicker of Lymph Node Metastases on 68Ga-PSMA Ligand PET/CT[J]. Clinical Nuclear Medicine, 2016, 41(10): 1
- [14] Machiels M, Wouterse S J, Geijsen E D, et al. Distribution of lymph node metastases on FDG-PET/CT in inoperable or unresectable oesophageal cancer patients and the impact on target volume definition in radiation therapy [J]. Journal of Medical Imaging & Radiation Oncology, 2016, 60(4): 520-527
- [15] Jilg C A, Drendel V, Beck T, et al. 563 The diagnostic accuracy of 68Ga-PSMA-PET/CT for detection of lymph node metastases in the setting of salvage lymph node dissection[J]. Journal of Urology, 2016, 195(4): e753-e753
- [16] Shigemoto Y, Suga K, Matsunaga N. F-18-FDG-avid lymph node metastasis along preferential lymphatic drainage pathways from the tumor-bearing lung lobe on F-18-FDG PET/CT in patients with non-small-cell lung cancer [J]. Annals of Nuclear Medicine, 2016, 30 (4): 287-297
- [17] Nakagawa M, Morimoto M, Takechi H, et al. Preoperative diagnosis of sentinel lymph node (SLN) metastasis using 3D CT lymphography (CTLG)[J]. Breast Cancer, 2016, 23(3): 519-524

- [18] Zhang Y, Luo Y K, Zhang M B, et al. Values of ultrasound features and MMP-9 of papillary thyroid carcinoma in predicting cervical lymph node metastases[J]. Scientific Reports, 2017, 7(1): 6670
- [19] Liang X, Yu J, Wen B, et al. MRI and FDG-PET/CT based assessment of axillary lymph node metastasis in early breast cancer: a meta-analysis[J]. Clinical Radiology, 2017, 72(4): 295-301
- [20] Khan S A, Zia S, Naqvi S U, et al. Relationship of Oral Tumor Thickness with the rate of lymph node metastasis in Neck based on CT Scan[J]. Pakistan Journal of Medical Sciences, 2017, 33(2): 353-357
- [21] Zhou L, Wang J Z, Wang J T, et al. Correlation analysis of MR/CT on colorectal cancer lymph node metastasis characteristics and prognosis [J]. Eur Rev Med Pharmacol Sci, 2017, 21(6): 1219-1225
- [22] Lin Z Y, Chen J. Treatment of recurrent mediastinal lymph node metastasis using CT-guided nontranspulmonary puncture interstitial implantation of (125)I seeds: Evaluation of initial effect and operative techniques[J]. Brachytherapy, 2016, 15(3): 361-369
- [23] Song B I, Kim H W, Won K S. Predictive Value of (18)F-FDG PET/CT for Axillary Lymph Node Metastasis in Invasive Ductal Breast Cancer[J]. Annals of Surgical Oncology, 2017, 24(8): 1-8
- [24] Kim J Y, Lee H, Kim T S, et al. Unusual Contralateral Axillary Lymph Node Metastasis in a Second Primary Breast Cancer Detected by FDG PET/CT and Lymphoscintigraphy [J]. Nuclear Medicine & Molecular Imaging, 2017, 15(Suppl 1): 1-4
- [25] Martínezesteve A, ÁlvarezPérez RM, Pachóngarrudo V M, et al. Mediastinal schwannoma mimicking lymph node metastasis from locally advanced breast cancer in a (18)F-FDG PET/CT study[J]. Revista Espanñola De Medicina Nuclear E Imagen Molecular, 2016, 35(4): 272
- [26] Wang Z L, Zhang X P, Tang L, et al. Lymph nodes metastasis of gastric cancer: Measurement with multidetector CT oblique multiplanar reformation-correlation with histopathologic results [J]. Medicine, 2016, 95(39): e5042
- [27] Atri M, Zhang Z, Dehdashti F, et al. Utility of PET-CT to Evaluate Retroperitoneal Lymph Node Metastasis in Advanced Cervical Cancer: Results of ACRIN6671/GOG0233 Trial [J]. Gynecologic Oncology, 2016, 2(3): 11-18
- [28] Chai Y, Gao J, Xing J, et al. Preoperative assessment value of spectral CT quantitative parameters in lymph node metastasis of gastric cancer [J]. Chinese journal of gastrointestinal surgery, 2017, 20(3): 309
- [29] Salman R, Hussain M, Adil S O. Diagnostic Accuracy of Multislice CT Scan in the Detection of Occult Cervical Lymph Node Metastasis in Head and Neck Cancers [J]. Journal of the College of Physicians and Surgeons-Pakistan: JCPSP, 2017, 27(5): 275
- [30] Hayashi H, Ashizawa K, Ogihara Y, et al. Comparison between solid component size on thin-section CT and pathologic lymph node metastasis and local invasion in T1 lung adenocarcinoma[J]. Japanese Journal of Radiology, 2017, 3(3): 1-7